

Alpha-7/MB™-1

Boehringer Ingelheim

Bacterin-Toxoid

Clostridium Chauvoei-Septicum-Novyi-Sordellii-Perfringens Types C & D-Moraxella Bovis Bacterin-Toxoid

U.S. Vet. Lic. No.: 124

**Indications:**

Recommended for the vaccination of healthy, susceptible cattle as an aid in the reduction of diseases caused by Clostridium chauvoei, Cl septicum, Cl novyi, Cl sordellii, Cl perfringens Types C and D, and for the prevention and control of pinkeye caused by Moraxella bovis.

Preservative: Gentamicin.

**Dosage and Administration:**

Inject a single 2 mL dose subcutaneously to cattle 2 months of age or older, prior to the onset of pinkeye season. Calves vaccinated under 3 months of age should be revaccinated at weaning or at 4-6 months of age. A 2 ml booster dose is recommended annually.

**Precautions:**

Store at a temperature of 35-45°F (2-7°C). Avoid freezing. Shake well before using. Use entire contents when first opened. Do not vaccinate within 60 days of slaughter. The use of a biological product in lactating dairy cattle can cause reduced milk production. Injection site swellings will often occur that may persist. Anaphylactoid reactions may occur.

Antidote: Epinephrine.

**Warnings:**

Animal inoculation only. Accidental injection to humans can cause serious local reactions. Contact a physician immediately if accidental injection occurs.

Presentation: 10 dose (20 mL) and 50 dose (100 mL) vials.