

Bar Vac® CD/T
Boehringer Ingelheim
Toxoid
Clostridium Perfringens Types C & D-Tetanus Toxoid
Composition: Prepared from cultures of the organisms listed. Alum precipitated.

Indications: Recommended for the vaccination of healthy, susceptible sheep, goats and cattle against enterotoxemia and tetanus caused by the toxins of Clostridium perfringens Types C and D and Clostridium tetani. Although Cl. perfringens Type B is not a significant problem in the U.S.A., immunity may be provided against the beta and epsilon toxins elaborated by Cl. perfringens Type B. This immunity is derived from the combination of Type C (beta) and Type D (epsilon) fractions.

Dosage and Administration:
Cattle: Using aseptic technique, inject 5 mL subcutaneously. Repeat in 21 to 28 days and once annually.
Sheep and Goats: Using aseptic technique, inject 2 mL subcutaneously. Repeat in 21 to 28 days and once annually.

Precaution(s): Store out of direct sunlight at 35-45°F (2-7°C). Avoid freezing. Shake well before using. Use entire contents when first opened.

CAUTION(s): Anaphylactoid reactions may occur.

ANTIDOTE(s): Administer epinephrine.

WARNING(s): Do not vaccinate within 21 days before slaughter.

Boehringer Ingelheim Vetmedica, Inc.
2621 North Belt Highway
St. Joseph, MO 64506-2002
1-800-325-9167
www.bi-vetmedica.com

Disclaimer

Every effort has been made by Stockyards to ensure the accuracy of the information listed above. However, it remains your responsibility to become familiar with the products you are purchasing. Please consult your family veterinarian if you have any questions.